

La rivoluzione dell'immunoterapia nel trattamento del melanoma

Margaret Ottaviano, MD, PhD

Unit of Melanoma, Cancer Immunotherapy and Innovative Therapies

*Istituto Nazionale Tumori – Fondazione "G. Pascale"
Director Paolo A. Ascierto*

DISCLOSURES

Speaker Fees, Travel and accommodation

Bristol-Myers-Squibb

MSD

Sun Pharma

Pierre Fabre

Novartis



Regeneron

Patient characteristics affecting immune surveillance

Active immune surveillance

Long-term benefit patients



- ≤ 3 brain metastases (size < 2 cm)
- Low tumour burden (< 3 organ involved?)
- Normal LDH



Inactive immune surveillance

No long-term benefit patients

- Multiple (>3) brain metastases
- High tumour burden (>3 organ involved?)
- High LDH



LDH, lactate dehydrogenase


Ascierto P, Dummer R. Oncoimmunology. 2018; Ascierto P, Ed. Session ASCO. 2019

Agenda

Immune checkpoint inhibitors

TIL therapy

Cancer vaccines




Oncolytic virus therapy

CAR-T cells

Agenda

Immune checkpoint inhibitors



TIL therapy

Oncolytic virus therapy

Cancer vaccines

CAR-T cells

Improved Survival with Ipilimumab in Patients with Metastatic Melanoma

The **NEW ENGLAND**
JOURNAL *of* **MEDICINE**

ESTABLISHED IN 1812


AUGUST 19, 2010

VOL. 363 NO. 8

F. Stephen Hodi, M.D., Steven J. O'Day, M.D., David F. McDermott, M.D., Robert W. Weber, M.D., Jeffrey A. Sosman, M.D., John B. Haanen, M.D., Rene Gonzalez, M.D., Caroline Robert, M.D., Ph.D., Dirk Schadendorf, M.D., Jessica C. Hassel, M.D., Wallace Akerley, M.D., Alfons J.M. van den Eertwegh, M.D., Ph.D., Jose Lutzky, M.D., Paul Lorigan, M.D., Julia M. Vaubel, M.D., Gerald P. Linette, M.D., Ph.D., David Hogg, M.D., Christian H. Ottensmeier, M.D., Ph.D., Celeste Lebbé, M.D., Christian Peschel, M.D., Ian Quirt, M.D., Joseph I. Clark, M.D., Jedd D. Wolchok, M.D., Ph.D., Jeffrey S. Weber, M.D., Ph.D., Jason Tian, Ph.D., Michael J. Yellin, M.D., Geoffrey M. Nichol, M.B., Ch.B., Axel Hoos, M.D., Ph.D., and Walter J. Urba, M.D., Ph.D.

— Ipi plus gp100 - - - Ipi - - - gp100
● ● ● Censored x x x Censored ■ ■ ■ Censored


A Overall Survival



No. at Risk

Ipi plus gp100	403	297	223	163	115	81	54	42	33	24	17	7	6	4	0
Ipi	137	106	79	56	38	30	24	18	13	13	8	5	2	1	0
gp100	136	93	58	32	23	17	16	7	5	5	3	1	0	0	0


B Progression-free Survival




No. at Risk

Ipi plus gp100	403	85	52	27	17	14	10	8	5	4	2	2	1	0
Ipi	137	37	26	17	13	10	10	9	6	4	2	1	0	0
gp100	136	18	7	5	3	2	2	2	1	0	0	0	0	0


Therapeutic revolution



Long-term OS in clinical trials with immuno-oncology agents and TT In advanced melanoma



F.L. 47 Yo




ORIGINAL ARTICLE

Final, 10-Year Outcomes with Nivolumab plus Ipilimumab in Advanced Melanoma


J.D. Wolchok, V. Chiarion-Sileni, P. Rutkowski, C.L. Cowey, D. Schadendorf, J. Wagstaff, P. Queirolo, R. Dummer, M.O. Butler, A.G. Hill, M.A. Postow, C. Gaudy-Marqueste, T. Medina, C.D. Lao, J. Walker, I. Márquez-Rodas, J.B.A.G. Haanen, M. Guidoboni, M. Maio, P. Schöffski, M.S. Carlino, S. Sandhu, C. Lebbé, P.A. Ascierto, G.V. Long, C. Ritchings, A. Nassar, M. Askelson, M.P. Benito, W. Wang, F.S. Hodi, and J. Larkin, for the CheckMate 067 Investigators*

CheckMate 067: study design and follow-up length



^aNo patient was being treated on study at the time of the final DBL. ^bFrom the date the last patient was randomized. ^cThe study was not powered for a comparison between NIVO + IPI and NIVO. ^dNIVO + IPI or NIVO alone vs IPI. ^eNIVO + IPI vs NIVO alone.


Overall survival



- At 10 years, NIVO + IPI and NIVO alone continued to demonstrate a significant and durable OS benefit

^aDescriptive comparison.

Melanoma-specific survival (MSS)^a




- After 10 years, median MSS was not reached (> 120 months) with NIVO + IPI and diverged from OS (52% vs 43%)
 - Results were consistent by BRAF mutation and PD-L1 expression status (Supplemental Figures S1 and S2)

^aIn this post hoc descriptive analysis, an event was defined as death due to melanoma or euthanasia due to disease progression; any other death was censored. ^bUnstratified. ^cDescriptive comparison.




OS and MSS in patients with PFS at 3 years


OS



MSS



- PFS at 3 years may be a good surrogate for long-term survival, with 10-year MSS rates $\geq 96\%$ in the NIVO-containing arms




December 2023:
4 cycle of combo ICIs
and 2 cycle of Nivolumab 480 mg...

BUT
Hepatitis G2 and
Colitis G2

RELATIVITY-047 is a global, randomized, double-blind, phase 2/3 study

Study design




RELATIVITY-047 (NCT03470922).

^aFirst tumor assessment (RECIST v1.1) was performed 12 weeks after randomization, every 8 weeks up to 52 weeks, and then every 12 weeks. ^bOS boundary for statistical significance was $P < 0.04302$ (2-sided) analyzed at 69% power; target HR, 0.75. ^cORR could not be formally tested and was descriptively analyzed. ^dLAG-3 expression on immune cells (1%) was determined by an analytically validated IHC assay (Labcorp, Burlington, NC, USA). ^ePD-L1 expression on tumor cells (1%) was determined by a validated Agilent Dako PD-L1 IHC 28-8 pharmDx test (Agilent, Santa Clara, CA, USA). ^fMinimum potential follow-up was defined as the time from last patient randomized to last patient, last visit.


RELATIVITY 047 demonstrated superior PFS benefit by BICR for RELA + NIVO vs NIVO

Figure 2. PFS per BICR



RELATIVITY 047 demonstrated superior OS benefit for RELA + NIVO vs NIVO

Figure 3. OS




Safety summary

	NIVO + IPI (n = 313)		NIVO (n = 313)		IPI (n = 311)	
	Any grade	Grade 3-4	Any grade	Grade 3-4	Any grade	Grade 3-4
Treatment-related AE, %	96	63	87	25	86	30
Treatment-related AE leading to discontinuation, %	45	34	16	9	17	15
Treatment-related death, ^a n (%)	2 (1)		1 (< 1)		1 (< 1)	

AE, n (%)	NIVO + RELA (n = 355)		NIVO (n = 359)	
	Any grade	Grade 3-4	Any grade	Grade 3-4
Any AE	352 (99)	164 (46)	345 (96)	141 (39)
TRAE	302 (85)	78 (22)	263 (73)	43 (12)
Leading to discontinuation	63 (18)	34 (10)	35 (10)	14 (4)
Treatment-related deaths ^a	4 (1)		2 (1)	


What can we do in future perspectives to avoid this ??

To implement screening campaigns




**To impact on earlier stages of disease
Neoadjuvant/Adjuvant In melanoma**

RFS curves in the current clinical practice of stage III Melanoma



Hauschild et al. ASCO 2020
Weber et al SMR 2021
Eggermont et al ESMO 2020


The problem with adjuvant therapy



- **ALL** get treated for 1 year
- Expensive (~\$100,000 pp)
- q3-4w clinic visits
- 3 monthly scans
- Toxicity
- No idea if treatment working


Finally a flat line!!!

Pooled analysis: Neoadjuvant Immunotherapy in stage III melanoma




Neoadjuvant immunotherapy for melanoma is now ready for clinical practice

Claus Garbe, Reinhard Dummer, Teresa Amaral, Rodabe N. Amaria, Paolo A. Ascierto, Elizabeth M. Burton, Brigitte Dreno, Alexander M. M. Eggermont, Axel Hauschild, Christoph Hoeller, Roland Kaufmann, Celeste Lebbe, Mario Mandala, Alexander M. Menzies, David Moreno, Olivier Michielin, Paul Nathan, Sapna P. Patel, Caroline Robert, Dirk Schadendorf, Paul C. Lorigan, Richard A. Scolyer, Hussein A. Tawbi, Bart A. van de Wiel, Christian Blank & Georgina V. Long





Individuals with cutaneous melanoma who are treated with neoadjuvant immunotherapy show a substantial improvement; this should be incorporated into standard care.

Neoadjuvant Therapy




Neoadjuvant-adjuvant phase 2 and 3 trials in macroscopic stage 3 melanoma SWOG 1801 and NADINA



No. at Risk

Neoadjuvant-adjuvant group	154	96	69	46	25	17	1
Adjuvant-only group	159	98	67	40	22	10	2




Patient 420 included November 30, 2023


Courtesy of Christian Blank

Patel et al. NEJM 2023

Distant metastasis-free survival



Updated event-free survival



Agenda


Immune checkpoint inhibitors

TIL therapy

Oncolytic virus therapy

Cancer vaccines

CAR-T cells



Why CAR-T cells?



CAR-T cell therapy for solid cancers

Clinical trial status


- Nearly 500 clinical trials evaluating CAR-T cells in solid tumors have been registered, many still ongoing
- Most completed studies are phase I/II trials that have reported modest results, with only occasional and generally brief clinical responses
- Clinical research on CAR-T cells has mainly focused on glioblastoma, sarcoma, neuroblastoma, and gastrointestinal cancer.

Albarran et al., Front Immunol 2024

Potential CAR-T cell targets for metastatic melanoma

Open access Original research Journal for Immunotherapy of Cancer Safety and biological outcomes following a phase 1 trial of GD2-specific CAR-T cells in patients with GD2-positive metastatic melanoma and other solid cancers

Tessa Gargett^{1,2,3} Nga T H Truong^{1,2} Bryan Gardam^{1,2} Wenbo Yu^{1,2} Lisa M Ebert^{1,2,3} Amy Johnson⁴ Erica C F Yeo¹ Nicole L Wittwer^{1,2} Gonzalo Tapia Rico^{2,3} Jesikah Logan² Purany Sivaloganathan² Maria Collis⁵ Andrew Ruszkiewicz^{2,3,5} Michael P Brown^{1,2,3}



5 MM patients with PR all treated with concomitant BRAFi/MEKi

2024;12:e008659. doi:10.1136/jitc-2023-008659

BARCELONA 2024 ESMO congress

Inge Marie Svane

nature communications



Article

<https://doi.org/10.1038/s41467-024-45221-2>

CAR-T cell therapy targeting surface expression of TYRP1 to treat cutaneous and rare melanoma subtypes


Received: 17 August 2023

Accepted: 18 January 2024

Published online: 09 February 2024

Check for updates

Sameeha Jilani^{1,†}, Justin D. Seco^{1,†}, Edume Mugerza^{1,2}, Aleida Pujol-Morillo¹, Jeffrey Chokry¹, Clement Ng¹, Gabriel Abili-Rodriguez^{1,2}, David Berger-Manerio¹, Ami Pant³, Jene Hu³, Rubi Gupta¹, Agustín Vega-Crespo¹, Ignacio Baselga-Carretero¹, Jia M. Chen^{1,2}, Daniel Sanghoo Shin^{4,5,6}, Philip Scumpia^{7,8}, Roxana A. Radu⁹, Yvonne Chen^{2,6,9,10}, Antoni Ribas^{1,2,6,10} & Cristina Puig-Saus^{1,2,6,10}



Antitumor activity in murine models on cutaneous, acral and uveal melanoma models with high TYRP1 overexpression

Content of this presentation is copyright and responsibility of the author. Permission is required for re-use.

Agenda


Immune checkpoint
inhibitors

TIL therapy

Oncolytic virus therapy

Cancer vaccines

CAR-T cells



Multicenter Double-Blind Phase 3 Trial of Canvaxin™ vs Placebo as Post Surgical Adjuvant in Metastatic Melanoma


Donald L. Morton, MD



N. Mozzillo, J. F. Thompson, M. C. Kelley, M. Faries,
S. Schneebaum, J. Wagner, E. Hersh, L. Schuchter,
M. Ross, D. Pertschuk, C. Nardo, R. Elashoff, G.
Gammon and the MMAIT-IV Clinical Trial Group




MMAIT-III Overall Survival (Intent To Treat)





MMAIT-IV Overall Survival (Intent To Treat)




Adjuvant Ganglioside GM2-KLH/QS-21 Vaccination Versus Observation After Resection of Primary Tumor > 1.5 mm in Patients With Stage II Melanoma: Results of the EORTC 18961 Randomized Phase III Trial

VOLUME 31 · NUMBER 30 · OCTOBER 20 2013

JOURNAL OF CLINICAL ONCOLOGY

ORIGINAL REPORT


Alexander M.M. Eggermont, Stefan Suciu, Piotr Rutkowski, Jeremy Marsden, Mario Santinami, Philippa Corrie, Steinar Aamdal, Paolo A. Ascierto, Poulam M. Patel, Wim H. Kruit, Lars Bastholt, Lorenzo Borgognoni, Maria Grazia Bernengo, Neville Davidson, Larissa Polders, Michel Praet, and Alan Spatz



No. at risk	O	n
Observation	204	657
GM2-KLH21	205	657

509	252	9
498	246	4

Negative!



No. at risk	O	n
Observation	112	657
GM2-KLH21	124	657

590	305	12
569	294	9



Individualized neoantigen therapy mRNA-4157 (V940) plus pembrolizumab in resected melanoma: 3-year update from the mRNA-4157-P201 (KEYNOTE-942) trial

Jeffrey S. Weber,¹ Muhammad Adnan Khattak,² Matteo S. Carlino,³ Tarek Meniawy,⁴ Matthew H. Taylor,⁵ George Anstas,⁶ Kevin B. Kim,⁷ Meredith McKean,⁸ Ryan J. Sullivan,⁹ Mark B. Faries,¹⁰ Thuy Tran,¹¹ C. Lance Cowey,¹² Theresa M. Medina,¹³ Jennifer M. Segar,¹⁴ Victoria Atkinson,¹⁵ Geoffrey T. Gibney,¹⁶ Jason J. Luke,¹⁷ Elizabeth I. Buchbinder,¹⁸ Georgina V. Long,¹⁹ INT Research and Development Author Group,^{20,21,a} Robert S. Meehan²⁰


^aManju Morrissey,²⁰ Igor Feldman,²⁰ Vasudha Sehgal,²⁰ Huzhang Mao,²⁰ Jia Guo,²⁰ Min Liu,²⁰ Anjali Rao,²⁰ Wei Zheng,²⁰ Praveen Aanur,²⁰ Lakshmi Srinivasan,²⁰ Mo Huang,²¹ Tal Zaks,²⁰ Michelle Brown,²⁰ Tracey Posadas²⁰

¹Laura and Isaac Perlmutter Cancer Center at NYU Langone Health, New York, NY, USA; ²Hollywood Private Hospital and Edith Cowan University, Perth, Australia; ³Melanoma Institute Australia and Westmead Hospital, Sydney, Australia; ⁴Saint John of God Subiaco Hospital, Subiaco, Australia; ⁵Earle A. Chiles Research Institute, Portland, OR, USA; ⁶Washington University School of Medicine, St Louis, MO, USA; ⁷California Pacific Medical Center Research Institute, San Francisco, CA, USA; ⁸Sarah Cannon Research Institute, Nashville, TN, USA; ⁹Massachusetts General Hospital, Boston, MA, USA; ¹⁰The Angeles Clinic and Research Institute, Los Angeles, CA, USA; ¹¹Yale-New Haven Hospital, New Haven, CT, USA; ¹²Baylor Charles A. Sammons Cancer Center, Dallas, TX, USA; ¹³University of Colorado, Aurora, CO, USA; ¹⁴University of Arizona Cancer Center, Tucson, AZ, USA; ¹⁵Princess Alexandra Hospital, Woolloongabba, Australia; ¹⁶Lombardi Comprehensive Cancer Center, Washington, DC, USA; ¹⁷UPMC Hillman Cancer Center, Pittsburgh, PA, USA; ¹⁸Dana-Farber Cancer Institute, Boston, MA, USA; ¹⁹Melanoma Institute Australia, Sydney, Australia; ²⁰Moderna, Inc., Cambridge, MA, USA; ²¹Merck & Co., Inc., Rahway, NJ, USA.

Sponsored by Moderna, Inc., in collaboration with Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA.

mRNA-4157-P201/KEYNOTE-942 (NCT03897881) study design

Randomized, phase 2, open-label study in patients with adjuvant resected melanoma at high risk of recurrence




Designed with 80% power to detect a hazard ratio of 0.5 with 40 RFS events (with a 1-sided alpha of 0.1 per protocol)
Primary analysis **triggered after a minimum of 1-year planned follow-up^c** (November 14, 2022 data cut) and at least 40 RFS events have been observed. DMFS analysis was prespecified for testing following positive RFS in the ITT population

Supportive analysis was **triggered after a minimum of 2 years of planned follow-up^c** (November 3, 2023 data cut)
Median planned follow-up^c: ~3yrs


^aPatients with stage IIIB disease were eligible only if relapse occurred within 3 months of prior surgery of curative intent; ^bAccording to the 8th edition of the American Joint Committee on Cancer Staging Manual ^cDefined as the time from the first dose date (or date of randomization if not treated) to date of clinical cut-off.
ECOG PS, Eastern Cooperative Oncology Group performance status; IM, intramuscular; ITT, intent-to-treat; IV, intravenous; NGS, next-generation sequencing; Q3W, every 3 weeks.

Sustained improvement of RFS primary efficacy endpoint



^aThe hazard ratio and 95% CI for mRNA-4157 (V940) + pembrolizumab versus pembrolizumab were estimated using a Cox proportional hazards model with treatment group as a covariate, stratified by disease stage (stages IIIB or IIIC or IIID vs stage IV) used for randomization. The *P* value is based on a 2-sided log-rank test stratified by disease stage (stages IIIB or IIIC or IIID vs stage IV) used for randomization; ^bFormal hypothesis testing of RFS was performed using November 2022 data cut. *P* value reported above used the November 2023 data cut; it's nominal and not for formal hypothesis testing. NE, not estimable.

Sustained improvement of DMFS secondary endpoint



^aThe hazard ratio and 95% CI for mRNA-4157 (V940) plus pembrolizumab versus pembrolizumab were estimated using a Cox proportional hazards model with treatment group as a covariate, stratified by disease stage (stages IIIB or IIIC or IIID vs stage IV) used for randomization. The *P* value is based on a 2-sided log-rank test stratified by disease stage (stages IIIB or IIIC or IIID vs stage IV) used for randomization; ^bFormal hypothesis testing of DMFS was performed using November 2022 data cut. *P* value reported above used the November 2023 data cut; it's nominal and not for formal hypothesis testing.

Personalized RNA neoantigen vaccines stimulate T cells in pancreatic cancer




Fig. 3 | mRNA vaccine response correlates with delayed PDAC recurrence. **a**, OS and RFS in $n = 19$ patients in the safety-evaluable cohort. **b**, RFS from surgery and from landmark time (date of the last vaccine priming dose) stratified by vaccine response in patients in the biomarker-evaluable cohort. n indicates individual patients. HR indicates hazard ratio with 95% CI. P values calculated using two-tailed log-rank test.




Fig. 4 | Vaccine-expanded T cells can infiltrate a micrometastasis. Clinical and immunological snapshot of a disappearing intrahepatic lymphoid aggregate after vaccination in a patient who responded to the vaccine. **a**, Serial percentage of vaccine-expanded T cells in blood analysed using CloneTrack and serum CA19-9 (left), and abdominal MRI (right) before and after vaccination. **b**, Haematoxylin and eosin staining (left), multiplexed immunofluorescence (middle) and percentage of vaccine-expanded T cells measured using CloneTrack (right, grey bar) in a new liver lesion that developed after vaccination detected

by MRI as in **a**. All 15 vaccine-expanded T cell clones (**a**, red line) were present in liver lesion (right, grey bar). **c**, Percentage of mutant TP53^{R175H} reads by digital droplet PCR in the liver lesion. The bar indicates the median, the error bars are the s.e.m. **d**, Uniform manifold approximation and projection (UMAP) plots of single-cell phenotypes of all blood T cells (left) and vaccine-expanded clones (middle), with effector markers (right). n indicates the number of T cells detected in liver lesion (**b**) or technical replicates (**c**). Data represent analyses of a single patient.

Agenda


Immune checkpoint
inhibitors

TIL therapy

Oncolytic virus therapy

Cancer vaccines

CAR-T cells



BACK
TO
THE **FUTURE**

TIL Therapy



Treatment with tumor-infiltrating lymphocytes (TIL) versus ipilimumab for advanced melanoma: results from a multicenter, randomized phase 3 trial

John B.A.G. Haanen, Maartje W. Rohaan, Troels Holz Borch, Joost H. van den Berg, Özcan Met, Marnix H. Geukes Foppen, Joachim Stoltenberg Granhøj, Bastiaan Nuijen, Cynthia Nijenhuis, Jos H. Beijnen, Inge Jedema, Maaïke van Zon, Inge Mansfield Noringriis, Rob Kessels, Sofie Wilgenhof, Johannes V. van Thienen, Ferry Lalezari, Alexander C.J. van Akkooi, Marco Donia, Inge Marie Svane

John B.A.G. Haanen


Paris, France, 10th September 2022

Presentation number LBA3




Tumor-infiltrating lymphocytes (TIL)

Preparation and treatment




Results (1)

Progression-free survival according to RECIST 1.1 in the ITT population




Results (4)

Overall survival in the ITT population



C-144-01 study design: autologous TIL therapy with lifileucel (LN-144) in advanced melanoma



Cohort 2 endpoints

- Primary: efficacy per investigator-assessed ORR using RECIST v1.1
- Secondary: safety and additional parameters of efficacy

Key eligibility criteria

- Radiographic confirmation of progression
- One tumor lesion resectable for TIL generation (~ 1.5 cm in diameter) and ≥ 1 target tumor lesion for RECIST v1.1 assessment
- Age ≥ 18 years at the time of consent
- ECOG PS 0-1

Methods

- Patients were enrolled from April 2017 to January 2019 at 26 sites across the US and EU
- Concomitant anticancer therapy was not permitted
- Imaging-evaluable disease was required
- All responses required confirmation
- Data cutoff: 22 April 2021


Adapted with permission from Larkin J.

BRAF, proto-oncogene B-Raf; Gen, generation; i, inhibitor; MEK, mitogen-activated protein kinase kinase; ORR, objective response rate; PD-1, programmed death 1; PFS, progression-free survival; PS, performance status; TIL, tumor-infiltrating leukocyte.

Larkin J et al. Presentation at the ASCO Annual Meeting; June 4-8, 2021; Virtual. Abstract 9505.

C-144-01: efficacy and safety

- 81% of patients had a reduction in tumor burden
- 17.7% of patients had further reduction of sum of diameters since the April 2020 data cutoff
- The ORR was 36.4% and the DCR was 80.3%
- Any-grade TEAEs were reported in 100% of patients
 - Grade 3-4: 97.0%
 - Grade 5: 3.0%



Adapted with permission from Larkin J.

This material may include information about investigational products and / or uses that are not approved for use in any country or in the country of your residence. Therefore, before prescribing any product, always refer to local materials such as the prescribing information and / or the Summary of Product Characteristics (SPC). Not all discussed therapies are approved for clinical use. Bristol Myers Squibb only recommends usage of approved products. Please check the product information of your country, approvals may vary. Refer to each country's local guidance for specific therapeutic strategies.

Median follow-up was 33.1 months (data cutoff, April 22, 2021). ^aPatients with BRAF V600 mutation. Three patients had no post-TIL disease assessments due to early death, and 1 due to start of new anticancer therapy.

BRAF, proto-oncogene B-Raf; CR, complete response; DCR, disease control rate; ORR, objective response rate; PD, progressive disease; PR, partial response; SD, stable disease; TIL, tumor-infiltrating leukocyte;

TEAE, treatment-emergent adverse event.

Larkin J et al. Presentation at the ASCO Annual Meeting; June 4-8, 2021; Virtual. Abstract 9505.

FDA NEWS RELEASE

FDA Approves First Cellular Therapy to Treat Patients with Unresectable or Metastatic Melanoma

For Immediate Release: February 16, 2024


Today, the U.S. Food and Drug Administration approved Amtagvi (lifileucel), the first cellular therapy indicated for the treatment of adult patients with a type of skin cancer (melanoma) that is unable to be removed with surgery (unresectable) or has spread to other parts of the body (metastatic) that previously has been treated with other therapies (a PD-1 blocking antibody, and if *BRAF* V600 mutation positive, a *BRAF* inhibitor with or without a MEK inhibitor).



IOV-COM-202: Phase 2, Multicohort, Multicenter Study of Lifileucel + Pembrolizumab in Patients With Solid Tumors


- Cohort 1A of IOV-COM-202 (NCT03645928) assesses the efficacy and safety of lifileucel + pembrolizumab in patients with ICI-naïve unresectable or metastatic melanoma
 - Patients may have received BRAF/MEK inhibitor treatment if they are *BRAF* mutation positive
 - Eligible patients must have ≥1 resectable lesion (≥1.5-cm diameter) and ≥1 measurable lesion for response assessment per RECIST v1.1
- Trial designed as a proof-of-concept study to support a registrational study in the frontline treatment setting

Treatment Schema



TILVANCE-301^a Global Phase 3 and Confirmatory Trial

Randomized study to evaluate lifileucel + pembrolizumab in frontline advanced melanoma
Enrolling in Europe, North America, and Australia



Study Endpoints

Dual primary efficacy endpoints

- BIRC-assessed ORR per RECIST v1.1
 - Potential for accelerated approval and confirmation of post anti-PD1 approval based on early interim analysis
- BIRC-assessed PFS per RECIST v1.1

Key secondary efficacy endpoint

- OS

Additional secondary endpoints

- BIRC-assessed CR rate, DOR, EFS per RECIST v1.1
- Investigator-assessed ORR, PFS, CR rate, DOR, EFS, PFS2 per RECIST v1.1
- Safety

^aNCT05727904.

BIRC, blinded independent review committee; CR, complete response; CY, cyclophosphamide; EFS, event-free survival; FLU, fludarabine; IL-2, interleukin-2; ORR, objective response rate; OS, overall survival; PD, progressive disease; PD-1, programmed cell death protein-1; PFS, progression-free survival; PFS2, progression-free survival 2; Q6W, every 6 weeks; RECIST, Response Evaluation Criteria in Solid Tumors.

Agenda


Immune checkpoint inhibitors

TIL therapy

Oncolytic virus therapy

Cancer vaccines

CAR-T cells




Primary efficacy, safety, and survival data from the registration-intended cohort of patients with anti-PD-1–failed melanoma from the IGNYTE clinical trial with RP1 combined with nivolumab

Caroline Robert¹, Mohammed M Milhem², Joseph J Sacco³, Judith Michels⁴, Gino K In⁵, Eva Muñoz Couselo⁶, Dirk Schadendorf⁷, Georgia M Beasley⁸, Jiaxin Niu⁹, Bartosz Chmielowski¹⁰, Trisha M Wise-Draper¹¹, Tawnya Lynn Bowles¹², Katy K Tsai¹³, Céleste Lebbé¹⁴, Caroline Gaudy-Marqueste¹⁵, Mark R Middleton¹⁶, Adel Samson¹⁷, Junhong Zhu¹⁸, Marcus Viana¹⁸, Michael K Wong¹⁹

¹Gustave Roussy and Paris-Saclay University, Villejuif, France; ²Holden Comprehensive Cancer Center, University of Iowa, Iowa City, IA, USA; ³The Clatterbridge Cancer Centre, Wirral, UK and University of Liverpool, Liverpool, UK; ⁴Département de Médecine Oncologique, Gustave Roussy, Villejuif, France; ⁵University of Southern California Norris Comprehensive Cancer Center, Los Angeles, CA, USA; ⁶Vall d'Hebron Institute of Oncology (VHIO) and Vall d'Hebron Hospital Medical Oncology Department, Barcelona, Spain; ⁷West German Cancer Center, University Hospital Essen, Essen, Germany; ⁸Duke Cancer Institute, Duke University, Durham, NC, USA; ⁹Banner MD Anderson Cancer Center, Gilbert, AZ, USA; ¹⁰Jonsson Comprehensive Cancer Center, University of California Los Angeles, Los Angeles, CA, USA; ¹¹University of Cincinnati Cancer Center, University of Cincinnati, Cincinnati, OH, USA; ¹²Intermountain Medical Center, Murray, UT, USA; ¹³Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, CA, USA; ¹⁴Université Paris Cité, AP-HP Dermato-Oncology and CIC, Cancer Institute APHP. Nord-Université Paris Cité, INSERM U976, Saint Louis Hospital, Paris, France; ¹⁵Aix-Marseille Université, APHM, Centre de Recherche en Cancérologie de Marseille (CRCM), INSERM, U1068, CNRS, UMR7258, UM105, Hôpital Timone, CEPCM, Dermatology and Skin Cancer Department, Marseille, France; ¹⁶Churchill Hospital and University of Oxford, Oxford, UK; ¹⁷Leeds Institute of Medical Research at St. James's, University of Leeds, Leeds, UK; ¹⁸Replimune, Inc., Woburn, MA, USA; ¹⁹University of Texas MD Anderson Cancer Center, Houston, TX, USA



Study design



Tumor response assessment: Radiographic imaging at baseline and every 8 weeks from first dose and every 12 weeks after confirmation of response

Primary objective

- Safety and efficacy using mRECIST* v1.1 by independent central review (sensitivity analysis by RECIST v1.1)

Secondary objectives

- ORR by investigator assessment (mRECIST* v1.1)
- DOR, CR rate, DOCB, DCR, and PFS by central and investigator assessment, 1-year and 2-year OS

* For mRECIST, PD must be confirmed by further progression at least 4 weeks after initial PD; intended to better allow for pseudoprogression than RECIST v1.1

Key eligibility

Anti-PD-1–failed advanced melanoma; measurable disease; adequate organ function; no prior oncolytic therapy; ECOG performance status 0–1

Criteria for prior anti-PD-1–failure

Confirmed progression while being treated with at least 8 weeks of anti-PD-1 therapy, alone or in combination; anti-PD-1 must be the last prior therapy. Patients on prior adjuvant therapy must have confirmed progression while being treated with adjuvant treatment (PD can be confirmed by biopsy)

Primary analysis conducted when all patients had ≥ 12 months follow-up

^aRP1 can be reinitiated beyond 8 cycles if protocol-specified criteria are met.

CR, complete response; DCR, disease control rate; DOCB, duration of clinical benefit; DOR, duration of response; ECOG, Eastern Cooperative Oncology Group; ORR, objective response rate; OS, overall survival; PD, progressive disease; PD-1, programmed cell death protein 1; PFS, progression-free survival; pfu, plaque-forming units; pt, patient; Q4W, every 4 weeks; RECIST, Response Evaluation Criteria in Solid Tumors.


Primary efficacy analysis

By blinded, independent central review

	Primary endpoint mRECIST v1.1 (N = 140)	Sensitivity analysis RECIST v1.1 (N = 140)
Confirmed best response, n (%)		
CR	21 (15.0)	21 (15.0)
PR	26 (18.6)	25 (17.9)
SD	41 (29.3)	31 (22.1)
PD	43 (30.7)	54 (38.6)
ORR (confirmed CR+PR), n (%)	47 (33.6)	46 (32.9)
95% CI	(25.8, 42.0)	(25.2, 41.3)

- 1 in 3 patients (33.6%) experienced a confirmed objective response, 15.0% CR

Overall survival



- One-, two-, and three-year survival rates were 75.3%, 63.3%, and 54.8%, respectively
- Median overall survival has not been reached

My personal Considerations

- Neoadjuvant combo ICI practice changing
- Still space for adjuvant therapy → Ongoing Phase III adjuvant trials are investigating in individualized neoantigen therapy
- IO COMBO as first line of treatment for advanced disease → something more to do for the safety profile
- Resistance to IO (BRAF wt) → ???????? (CAR-T?, Oncolytic virus)
- TILs in front line setting...

Paolo Antonio Ascierto

Patients, Families and Caregivers



ISTITUTO NAZIONALE TUMORI
IRCCS - Fondazione Pascale

Medical Oncologists

Ester Simeone
Lucia Festino
Vito Vanella
Claudia Trojaniello
Maria Grazia Vitale
Margaret Ottaviano
Francesca Sparano
Arianna Facchini

Biologists

Mariaelena Capone
Gabriele Madonna
Domenico Mallardo
Marilena Tuffanelli
Marilena Romanelli

Dermatologists

Rossella Di Trollo
Lugi Scarpato
Marco Palla
Mirella D'Andrea
Elisa Camela

Nurses

Patrizia Sabatelli
Federica Hauber
M Grazia Capasso
Giuseppina Marano
Teresa De Cristofaro
Angela Mazzocchi
Assunta Calignano

Study Coordinators

Assunta Esposito
Miriam Paone
Gianni Rinaldi
Benedetta Alfano
Mario Mallardo
Maria Rosaria Cecco



Margaret.Ottaviano@istitutotumori.na.it